

1. Find the gradient vector for the scalar function. (That is, find the conservative vector field for the potential function.)

$$f(x, y) = 7x^2 + 8xy + 2y^2$$

2. Find the gradient vector for the scalar function. (That is, find the conservative vector field for the potential function.)

$$f(x, y) = \sin 9x \cos 3y$$

3. Determine whether the vector field is conservative.

$$\vec{F}(x, y) = 9y^2(7y\hat{i} - x\hat{j})$$

- A) Conservative  
B) Not Conservative

4. Determine whether the vector field is conservative. If it is, find a potential function for the vector field.

$$\vec{F}(x, y) = 7x^6y\hat{i} + x^7\hat{j}$$

5. Determine whether the vector field is conservative. If it is, find a potential function for the vector field.

$$\vec{F}(x, y) = \frac{6y}{x}\hat{i} - \frac{x^6}{y^6}\hat{j}$$

6. Find the curl for the vector field at the given point.

$$\vec{F}(x, y, z) = 2xyz\hat{i} + 2y\hat{j} + 2z\hat{k}, \quad (2, 3, 2)$$

7. Determine whether the vector field is conservative. If it is, find a potential function for the vector field.

$$\vec{F}(x, y, z) = 3x^2y^4z^5\hat{i} + 4x^3y^3z^5\hat{j} + 5x^3y^4z^4\hat{k}$$

8. Find the divergence of the vector field.

$$\vec{F}(x, y, z) = 9x^4\hat{i} - xy^3\hat{j}$$

9. Find the divergence of the vector field at the given point.

$$\vec{F}(x, y, z) = 8xyz\hat{i} + 8y\hat{j} + 8x\hat{k}, \quad (8, 9, 8)$$

10. **OMIT**

11. **OMIT**

12. Evaluate the line integral along the given path.

$$\int_C (2x - 7y) ds \quad C: \vec{r}(t) = 7t\hat{i} + 3t\hat{j}, \quad 0 \leq t \leq 8$$

13. Evaluate  $\int_C (x^2 + y^2) ds$  where the path  $C$  is:

(i) the  $x$ -axis from  $x = 0$  to  $x = 1$ ;

(ii) the  $y$ -axis from  $y = 1$  to  $y = 3$ .

14. Evaluate  $\int_C (x + 49\sqrt{y}) ds$  where the path  $C$  is:

(i) the line from  $(0,0)$  to  $(1,1)$ ;

(ii) the line from  $(0,0)$  to  $(3,9)$ .

15. Evaluate  $\int_C \vec{F} \cdot d\vec{r}$  where  $C$  is represented by  $\hat{r}(t)$ .

$$\vec{F}(x, y) = xy\hat{i} + y\hat{j}$$

$$C: \vec{r}(t) = 32t\hat{i} + t\hat{j}, \quad 0 \leq t \leq 8$$

16. Evaluate  $\int_C \vec{F} \cdot d\vec{r}$  where  $C$  is represented by  $\vec{r}(t)$ .

$$\vec{F}(x, y) = 5x\hat{i} + 9y\hat{j}$$

$$C: \vec{r}(t) = t\hat{i} + \sqrt{4-t^2}\hat{j}, \quad -2 \leq t \leq 2$$

17. Evaluate the line integral along the path  $C$  given by  $x = 4t$ ,  $y = 20t$ , where  $0 \leq t \leq 1$ .

$$\int_C (x + 4y^2) ds$$

18. Evaluate the line integral

$$\int_C (2x - y) dx + (x + 3y) dy$$

along the path  $C$ , where  $C$  is:

(i) the  $x$ -axis from  $x = 0$  to  $x = 8$ .

(ii) the  $y$ -axis from  $y = 0$  to  $y = 12$ .

19. Find the area of the lateral surface over the curve  $C$  in the  $xy$ -plane and under the surface  $z = f(x, y)$ , where

$$\text{Lateral surface area} = \int_C f(x, y) ds$$

$f(x, y) = 4$ ,  $C$ : line from  $(0, 0)$  to  $(5, 6)$ .

20. Set up and evaluate the integral  $\int_C \vec{F} \cdot d\vec{r}$  for each parametric representation of  $C$ .

$$\vec{F}(x, y) = x^2 \hat{i} + xy \hat{j}$$

(i)  $\vec{r}_1(t) = 3t \hat{i} + 7t^2 \hat{j}$ ,  $0 \leq t \leq 1$

(ii)  $\vec{r}_2(\theta) = 3 \sin \theta \hat{i} + 7 \sin^2 \theta \hat{j}$ ,  $0 \leq \theta \leq \frac{\pi}{2}$

21. Determine whether or not the vector field is conservative.

$$\vec{F}(x, y) = 40x^7 y^7 \hat{i} + 35x^8 y^6 \hat{j}$$

A) Conservative

B) Not conservative

22. Find the value of the line integral  $\int_C \vec{F} \cdot d\vec{r}$ , where  $\vec{F}(x, y) = 2xy \hat{i} + x^2 \hat{j}$ .

(i)  $\vec{r}_1(t) = t \hat{i} + t^3 \hat{j}$ ,  $0 \leq t \leq 1$

(ii)  $\vec{r}_2(t) = t \hat{i} + t^7 \hat{j}$ ,  $0 \leq t \leq 1$

23. Find the value of the line integral  $\int_C (2x - 5y + 7) dx - (5x + 7y - 8) dy$ .

(i)  $C$ : the curve  $x = \sqrt{49 - y^2}$  from  $(0, -7)$  to  $(0, 7)$

(ii)  $C$ : from  $(0, -7)$  to  $(0, 7)$  along  $x = \sqrt{49 - y^2}$ , then back to  $(0, -7)$  along the  $y$ -axis (a closed curve).

24. Find the value of the line integral  $\int_C \vec{F} \cdot d\vec{r}$ , where  $\vec{F}(x, y) = 3yz \hat{i} + 3xz \hat{j} + 3xy \hat{k}$ .

(i)  $\vec{r}_1(t) = t \hat{i} + 9 \hat{j} + t \hat{k}$ ,  $0 \leq t \leq 81$

(ii)  $\vec{r}_2(t) = t^2 \hat{i} + 9 \hat{j} + t^2 \hat{k}$ ,  $0 \leq t \leq 9$

25. Evaluate the line integral using the Fundamental Theorem of Line Integrals. Use a computer algebra system to verify your results.

$$\int_C (4y\hat{\mathbf{i}} + 4x\hat{\mathbf{j}}) \cdot d\vec{\mathbf{r}}$$

$C$ : a smooth curve from  $(0,0)$  to  $(9,4)$

26. Evaluate the line integral using the Fundamental Theorem of Line Integrals. Use a computer algebra system to verify your results.

$$\int_C \frac{2x}{(x^2 + y^2)^2} dx + \frac{2y}{(x^2 + y^2)^2} dy$$

$C$ : circle  $(x-6)^2 + (y-3)^2 = 49$  clockwise from  $(13,3)$  to  $(-1,3)$

27. **OMIT**

28. Use Green's Theorem to evaluate the integral

$$\int_C (y-x)dx + (5x-y)dy$$

for the path  $C$ : boundary of the region lying between the graphs of  $y = x$  and  $y = x^2 - 18x$

29. Use Green's Theorem to evaluate the integral

$$\int_C 13xydx + (x+y)dy$$

for the path  $C$ : boundary of the region lying between the graphs of  $y = 0$  and  $y = 169 - x^2$ .

30. Use Green's Theorem to evaluate the integral

$$\int_C (x^2 - y^2) dx + 10xy dy$$

for the path  $C: x^2 + y^2 = 9$ .

31. Use Green's Theorem to evaluate the integral

$$\int_C 8xy dx + 8(x + y) dy$$

for  $C$ : boundary of the region lying between the graphs of  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 16$ .

32. Use Green's Theorem to calculate the work done by the force  $\vec{F}$  on a particle that is moving counterclockwise around the closed path  $C$ .

$$\vec{F}(x, y) = 4xy\hat{i} + (x + y)\hat{j}$$

$$C: x^2 + y^2 = 9$$

33. Find the rectangular equation for the surface by eliminating parameters from the vector-valued function. Identify the surface.

$$\vec{r}(u, v) = u\hat{i} + v\hat{j} + \frac{v}{9}\hat{k}$$

34. Find a vector-valued function whose graph is the cylinder  $x^2 + y^2 = 4$ .

35. Find a vector-valued function whose graph is the ellipsoid  $\frac{x^2}{49} + \frac{y^2}{100} + \frac{z^2}{81} = 1$ .

36. Write a set of parametric equations for the surface of revolution obtained by revolving the graph of the function about the given axis.

$$y = \frac{x}{7}, \quad 0 \leq x \leq 21 \quad x\text{-axis}$$

37. Find an equation of the tangent plane to the surface represented by the vector-valued function at the given point.

$$\vec{r}(u, v) = (10u + v)\hat{i} + (u - v)\hat{j} + v\hat{k}, \quad (6, -6, 6)$$

38. Find the area of the surface over the given region. Use a computer algebra system to verify your results.

The sphere,

$$\vec{r}(u, v) = 8\sin u \cos v\hat{i} + 8\sin u \sin v\hat{j} + 8\cos u\hat{k}, \quad 0 \leq u \leq \pi, \quad 0 \leq v \leq 2\pi$$

39. Find the area of the surface over the given region. Use a computer algebra system to verify your results.

The part of the cone,

$$\vec{r}(u, v) = 3u \cos v\hat{i} + 3u \sin v\hat{j} + u\hat{k}$$

where  $0 \leq u \leq 8$  and  $0 \leq v \leq 2\pi$ .

40. Evaluate  $\iint_S (x - 7y + z) dS$ , where

$$S: z = 14 - x, \quad 0 \leq x \leq 14, \quad 0 \leq y \leq 14$$

41. Evaluate  $\iint_S f(x, y) dS$ , where

$$f(x, y) = y + 6$$

$$S: r(u, v) = u\hat{i} + v\hat{j} + \frac{v}{10}\hat{k}, \quad 0 \leq u \leq 1, \quad 0 \leq v \leq 10$$

42. Evaluate  $\iint_S f(x, y, z) dS$ , where

$$f(x, y, z) = x^2 + y^2 + z^2$$

$$S: x^2 + y^2 = 4, \quad 0 \leq z \leq 4$$